

	Category	Significant anticholinergic activity*	Moderate anticholinergic activity†	Mild anticholinergic activity‡
Anticholinergic drugs	Antiemetic/anti-vertigo	dimenhydrinate [§] , promethazine, prochlorperazine		metoclopramide
	Anti-parkinson	benzhexol, benztropine, biperiden, orphenadrine	amantadine	
	Antispasmodics (gastrointestinal)	atropine, hyoscine, propantheline	belladonna alkaloids [§]	
	Antispasmodics (genitourinary)	oxybutynin, darifenacin, solifenacin, tolterodine		
	Bronchodilators		ipratropium, tiotropium	
	Mydriatics/cycloplegics	atropine, cyclopentolate, homatropine, tropicamide		
Drugs with anticholinergic side effects	Analgesics	pizotifen	pethidine	codeine, colchicine, fentanyl, morphine, oxycodone
	Antiarrhythmics			atenolol, digoxin, disopyramide, metoprolol
	Antidepressants	amitriptyline, clomipramine, dothiepin, doxepin, imipramine, nortriptyline, paroxetine, trimipramine	mirtazapine	bupropion, fluoxetine, fluvoxamine
	Antidiarrhoeals	diphenoxylate, atropine [§]		loperamide
	Antihistamines	chlorpheniramine [§] , dexchlorpheniramine, dimenhydrinate [§] , diphenhydramine [§] , promethazine	cyproheptadine	brompheniramine maleate [§]
	Antipsychotics	chlorpromazine, clozapine, olanzapine, pericyazine, quetiapine, thioridazine, trifluoperazine	pimozide	haloperidol, risperidone
	Bronchodilators			theophylline
	Miscellaneous		baclofen, carbamazepine, lithium, pseudoephedrine	captopril, chlorthalidone, dipyridamole, frusemide, hydralazine, isosorbide, nifedipine, prednisolone, warfarin

* Drugs with clinically significant anticholinergic activity.

† Drugs noted to have anticholinergic adverse effects, usually at excessive doses.

‡ Drugs with potential anticholinergic activity (according to receptor binding).

§ Also part of multi-ingredient preparations available over the counter.

References

- Mintzer J, et al. Anticholinergic side-effects of drugs in elderly people. *J R Soc Med* 2000;93:457–62.
- Boustani M, et al. Impact of anticholinergics on the aging brain: a review and practical application. *Aging Health* 2008;4:311–20.
- Rudolph J, et al. The anticholinergic risk scale and anticholinergic adverse effects in older persons. *Arch Intern Med* 2008;168:508–13.
- Chew M, et al. Anticholinergic activity of 107 medications commonly used by older adults. *J Am Geriatr Soc* 2008;56:1333–41.
- Carnahan R, et al. The anticholinergic drug scale as a measure of drug-related anticholinergic burden: associations with serum anticholinergic activity. *J Clin Pharmacol* 2006;46:1481–86.

NPS is an independent, non-profit organisation for Quality Use of Medicines, funded by the Australian Government Department of Health and Ageing.